European Lobster (Homarus gammarus) Import to Iceland



Ragnheidur Thorarinsdottir Soffia Karen Magnusdottir Halldor Palmar Halldorsson

Objectives

- Import European lobster juveniles to Iceland from UK and Norway
- Perform growth and survival tests in two different setups with flow-through of very clean borehole water and semi-RAS using surface seawater
- Monitor growth at different temperatures and follow environmental parameters
- Test commercially available Arctic char feed as it has a very similar overall combination as the lobster feed developed by NLF
- Respiration tests

Lobster transport



Lobsters from UK:

(Age 6 weeks)
Arrived in circular trays in plastic bags with added oxygen

Lobsters from IMR:

(Age 22 months)
Arrived in a styrofoam box,
wrapped in moist newspaper



Experimental setup





Sudurnes Science and Learning Center			Saebyli
Drill h	Drill hole sea water, flow through		Surface sea water, semi RAS
	ters from way	NLH lobster from UK	NLH lobster from UK
10°C	16°C	16°C	17°C










Environment





T:10°C and 16/17°C

Salinity: 33 ppt


DO: >91%

CO2: 13.7-16.2 mg/L











formities



















Respiration tests









norden

Nordic Innovation











Technical University of Denmark







http://eurolobster.svinna.is/

